

WHAT VACCINE DOES AAACE RECOMMEND FOR ADULTS WITH DIABETES

CDC-Recommended Vaccines	When and How Often
Age-appropriate vaccines	<ul style="list-style-type: none"> All persons should receive according to the CDC schedules
COVID-19	<ul style="list-style-type: none"> One or more doses per current CDC recommendations and FDA approvals
Flu	<ul style="list-style-type: none"> All adults, annually
Hepatitis B	<ul style="list-style-type: none"> Adults aged 19-59 years Adults aged 60 years and older with risk factors for Hepatitis B 2, 3, or 4 doses depending on vaccine or condition
Pneumococcal - to protect against pneumonia	<ul style="list-style-type: none"> Adults with diabetes ages ≥ 19 years One or two doses, depending upon the type of vaccine received
RZV (Zoster) - to protect against shingles	<ul style="list-style-type: none"> All adults ≥ 50 years Adults ≥ 19 years who have weakened immune systems Two doses
Tdap - to protect against tetanus, diphtheria, and whooping cough	<ul style="list-style-type: none"> Every 10 years after first series (may need additional doses during pregnancy or after an injury)
RSV - to protect against respiratory syncytial virus	<ul style="list-style-type: none"> Adults aged 60 years or older who are at increased risk for severe RSV disease including those with chronic medical conditions such as diabetes

ACIP = Advisory Committee on Immunization Practices; CDC = Centers for Disease Control and Prevention

For information on the most current CDC vaccine recommendations for adults with diabetes, visit <https://www.cdc.gov/vaccines/hcp/imz-schedules/adult-schedule-vaccines.html>

